

RE-FEEDING **SEVERELY UNDERWEIGHT** **DONKEYS**

Re-feeding should be a gradual process and can take a minimum of 60-90 days for the transition back to normal body weight.

 11 October 2017  [Read on The Donkey Sanctuary website](#)

There are no specific guidelines regarding feeding severely underweight donkeys (BCS < 1.5) so current guidelines are based on recommendations made by Brian Walldridge (Kentucky Equine Research) on feeding malnourished horses, as well as our experience of such cases here at The Donkey Sanctuary.

Caution should be taken with the initial period of re-feeding as early overfeeding can result in refeeding syndrome where neurological signs are sometimes apparent and even death can occur due to cardiac or pulmonary failure.

Re-feeding should be a gradual process and can take a minimum of 60-90 days for the transition back to normal body weight. For a severely starved animal it may take as long as 6-10 months.

RE-FEEDING DONKEY **TIPS**

- Re-feeding should be very gradual and kept simple initially with only good quality hay or high fibre haylage (e.g. Horsehage High Fibre) being fed (unless the donkey has dental problems and cannot manage long fibre).
- Donkeys should have access to grazing (care should be taken not to turn out on lush or frosty grazing), initially for periods of about 2-3 hours at a time, increasing this by an hour every 2-3 days.
- Donkeys should have access to a vitamin and mineral block as a minimum but ideally should be fed a balancer such as Top Spec Comprehensive gradually building up to be fed at a rate of 100g/100Kg BW.

- Donkeys should be kept warm by the provision of **shelter** and **rugs**.
- Thin donkeys with long coats may need clipping in summer as they are likely to be more susceptible to heat stress as their metabolic rate rises

SUGGESTED **RE-FEEDING PROGRAMME**

DAY 0-3

Initially donkeys should be fed at 50% of their current (poor body condition) energy requirements (0.75% of BW in dry matter). Good quality hay or high fibre haylage (e.g. Horsehage High Fibre) should be all that is fed initially, fed in small quantities (up to $\frac{1}{4}$ Kg) every 4-6 hours. If teeth are poor a low energy and low non-structural carbohydrate (NSC) hay replacement product (e.g. Happy Hoof) can be fed instead.

DAY 4

After the initial 2-3 day adaptation period the donkey can be fed to 100% of its current weight requirements (up to $\frac{1}{2}$ kg) every 4-6 hours.

DAY 6-14

Over the next week or so the amount of hay fed can be increased to that of the donkey's ideal bodyweight (1.5% of BW in dry matter).

DAY 15 ONWARDS

Allow the donkey adlib access to forage, this may be good quality hay, high fibre haylage or a combination of the two. For those with poor teeth a hay replacement product can be fed at the rate of 2-3kg per day.

At this point it would be a good idea to introduce a feed balancer such as TopSpec Comprehensive which, in addition to vitamins and minerals, will provide the donkey with a moderate level of protein to help rebuild lost muscle.

Once the donkey is approaching a body condition score of 2 it would be wise to start introducing straw back into the diet in anticipation of a more normal straw based diet for the future.

Once the donkey has reached a body condition score of 3 and is otherwise healthy, the balancer can be changed to TopSpec Donkey Forage Balancer which is more appropriate for maintenance. Some donkeys may benefit from additional supplementary nutritional support during the refeeding period.

NOTE

It is important to bear in mind that an animal that has been unable to stand for 72 hours or more will have a very poor prognosis for recovery, even with aggressive treatment, as will animals that have lost 45-50% of their bodyweight.

MORE INFORMATION

Our skilled and experienced team of welfare advisers are on hand to give advice and appropriate support during office hours. For feeding and nutrition advice [contact our nutrition team](#) or telephone 01395 578222 and ask to be put through to the welfare department.